

SPECIFICATION

CITALOPRAM HYDROBROMIDE CRYSTAL AND METHOD FOR CRYSTALLIZATION THEREOF

TECHNICAL FIELD OF THE INVENTION

5 The present invention relates to a method for crystallizing citalopram hydrobromide useful as an antidepressant and citalopram hydrobromide having specific crystal properties.

BACKGROUND OF THE INVENTION

Citalopram hydrobromide is useful as an antidepressant.
10 When it is used as a pharmaceutical bulk for the preparation of a pharmaceutical agent, particle size, particle size distribution and aspect ratio of the crystal are the important crystal characteristics. However, there is no teaching as to what crystal characteristics of citalopram hydrobromide are required as a
15 beneficial pharmaceutical bulk, or a method for achieving such beneficial crystal characteristics.

According to US4650884, for example, a method for crystallizing citalopram hydrobromide comprises, crystallization in a single solvent of acetone, first recrystallization in a
20 single solvent of water, second recrystallization in a mixed solvent of methanol and isopropyl alcohol, and third recrystallization in a mixed solvent of methanol, acetone and hexane. This method aims at purification of citalopram crystals but not at controlling particle size, particle size distribution
25 or aspect ratio of crystals, not to mention a method for controlling these.

When citalopram hydrobromide is crystallized according to the method described in this publication, fine crystals having a particle size of less than 5 μm are formed in a large amount
30 (41.9% of the total amount). When fine crystals are contained in a large amount, filtering performance after crystallization for medicament production is degraded. In addition, the fine particles scatter and the workers are exposed thereto during production of pharmaceutical preparations, which is an
35 unpreferable aspect as a pharmaceutical bulk.

When citalopram hydrobromide is crystallized according to the method described in this publication, crystals having a smaller average aspect ratio of less than 2 are produced. Such

crystals cause problems such as poor filtering performance after crystallization for the preparation of a pharmaceutical agent and poor fluidity when crystals are being taken out, which is not preferable as a pharmaceutical bulk.

5 It is therefore an object of the present invention to provide

(1) a method for industrially crystallizing citalopram hydrobromide, which method affords easy control of crystal characteristics, such as particle size, particle size distribution
10 and aspect ratio of the crystals, and

(2) citalopram hydrobromide crystals having crystal characteristics useful as a pharmaceutical bulk.

SUMMARY OF THE INVENTION

Such object can be achieved by the present invention.
15 According to the present invention, citalopram hydrobromide dissolved in a specific solvent is crystallized or recrystallized while controlling the cooling rate, thereby to control the particle size, particle size distribution and aspect ratio of citalopram hydrobromide crystals. It has been also found that
20 citalopram hydrobromide crystals containing crystals having a particle size of less than of 5 μm in a proportion of 35% at most and citalopram hydrobromide crystals having an average aspect ratio of not less than 4.5 and not more than 6.0, which are obtained by this method, are advantageous as a pharmaceutical bulk.

25 Accordingly, the present invention provides the following.

(1) Citalopram hydrobromide crystals containing crystals having a particle size of less than 5 μm in a proportion of 35% at most.

(2) Citalopram hydrobromide crystals containing crystals having a particle size of less than 5 μm in a proportion of 35% at most,
30 and containing crystals having a particle size of not less than 20 μm in a proportion of not less than 10%, or having an average aspect ratio of not less than 2.0 and not more than 9.0, or having an average aspect ratio of not less than 2.5 and less than 4.5, or
35 than 6.0.

(3) Citalopram hydrobromide crystals having an average aspect ratio of not less than 2.0 and not more than 9.0.

(4) Citalopram hydrobromide crystals having an average aspect ratio of not less than 2.5 and less than 4.5.

(5) Citalopram hydrobromide crystals having an average aspect ratio of not less than 4.5 and not more than 6.0.

5 (6) A method for crystallizing citalopram hydrobromide, which comprises the steps of

(A1) dissolving, by heating, citalopram hydrobromide in a solvent containing at least one member selected from the group consisting of alcohol having 1 to 3 carbon atoms, water and acetone and

10 (B1) cooling the resulting product to achieve crystallization, while controlling the cooling rate.

(7) A method for crystallizing citalopram hydrobromide, which comprises the steps of

15 (A2) dissolving, by heating, citalopram hydrobromide in a solvent containing at least one member selected from the group consisting of alcohol having 1 to 3 carbon atoms, water and acetone,

(B2) cooling the obtained solution to achieve crystallization,

(C2) dissolving a part of the obtained crystals by heating, and

20 (D2) recrystallizing the crystals while controlling the cooling rate.

A preferable mode of the above-mentioned (6) includes

(6-1) controlling the cooling rate within the temperature range of 0°C - 80°C,

25 (6-2) controlling the average cooling rate within the temperature range of 20°C - 40°C to not less than 30°C/hour and not more than 60°C/hour,

(6-3) controlling the average cooling rate within the temperature range of 20°C - 40°C to not less than 0.5°C/hour and less than 30°C/hour, and

30 (6-4) cooling the solution obtained in (A1) to a temperature range of not less than 30°C and less than 48°C and adding a seed crystal of citalopram hydrobromide to achieve crystallization.

A preferable mode of the above-mentioned (7) includes

35 (7-1) cooling to the temperature range of not less than 30°C and less than 48°C in (B2),

(7-2) after cooling to the temperature range of not less than 30°C and less than 48°C, adding a seed crystal of citalopram hydrobromide to achieve crystallization in (B2),

(7-3) heating to the temperature range of not less than 48°C and not more than 60°C to dissolve a part of the crystal in (C2), (7-4) controlling the average cooling rate in the temperature range of from (heating temperature in (C2)) to (this heating temperature - 30°C) to not less than 30°C/hour and not more than 90°C/hour in (D2), and (7-5) controlling the average cooling rate in the temperature range of from (heating temperature in (C2)) to (this heating temperature - 30°C) to not less than 1°C/hour and less than 30°C/hour in (D2).

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows ¹H-NMR of the compound obtained in Reference Example 1.

Figure 2 shows a powder X-ray crystal diffraction pattern of the compound obtained in Reference Example 1.

Figure 3 shows a powder X-ray crystal diffraction pattern of the compound obtained in Example 1.

Figure 4 shows a powder X-ray crystal diffraction pattern of the compound obtained in Example 4.

Figure 5 shows a powder X-ray crystal diffraction pattern of the compound obtained in Comparative Example 1.

Figure 6 shows a crystal particle size distribution of the compound obtained in Example 1.

Figure 7 shows a crystal particle size distribution of the compound obtained in Example 2.

Figure 8 shows a crystal particle size distribution of the compound obtained in Example 3.

Figure 9 shows a crystal particle size distribution of the compound obtained in Example 4.

Figure 10 shows a crystal particle size distribution of the compound obtained in Example 5.

Figure 11 shows a crystal particle size distribution of the compound obtained in Example 6.

Figure 12 shows a crystal particle size distribution of the compound obtained in Comparative Example 1.

Figure 13 shows a micrograph (400 magnifications) of the compound obtained in Example 1.

Figure 14 shows a micrograph (400 magnifications) of the

compound obtained in Example 4.

Figure 15 shows a micrograph (400 magnifications) of the compound obtained in Example 6.

Figure 16 shows a micrograph (400 magnifications) of the compound obtained in Comparative Example 1.

Figure 17 shows a micrograph (400 magnifications) of a ruler graduated in 10 μm .

DETAILED DESCRIPTION OF THE INVENTION

The present invention is explained in detail in the following.

The average particle size in the present invention is an average diameter upon conversion of the volume of the crystal particles into sphere.

The average aspect ratio in the present invention is an average value of the major axis/minor axis of crystal particles.

According to the present invention, crystal characteristics such as particle size, particle size distribution and aspect ratio of the crystals can be controlled easily. As the industrial method for crystallizing citalopram hydrobromide includes the following methods.

Method 1

A first method for crystallizing citalopram hydrobromide in the present invention includes

(A1) dissolving citalopram hydrobromide in a specific solvent, and

(B1) crystallization by cooling the resulting solution while controlling a cooling rate (hereinafter this method is to be referred to as Method 1). The crystals obtained according to Method 1 generally show a smooth cone-shaped particle size distribution.

In the following, the steps (A1) and (B1) are explained in detail. In each step, it is preferable that the particles be stirred at 10 rpm - 250 rpm, particularly at 50 rpm - 220 rpm, so that the temperature of the system as a whole can be uniform and the crystals after stirring will not have a smaller aspect ratio.

According to Method 1, the cooling rate of the solution within the specific temperature range should be regulated to control the crystal characteristics. That is, the present

inventors have found that the cooling rate of the solution within the temperature range of preferably 0°C - 80°C, particularly 20°C - 40°C, in (B1) affects the crystal particle size of citalopram hydrobromide. For example, the crystals obtained when the average cooling rate of the solution within the temperature range of 20°C - 40°C is controlled to not less than 0.5°C/hour and less than 30°C/hour contain crystals having a particle size of less than 5 µm in a proportion of 35% at most, and have an average particle size of about 10 µm - 20 µm, and that the crystals obtained when it is controlled to not less than 30°C/hour and not more than 60°C/hour contain crystals having a particle size of less than 5 µm in a proportion of 35% at most, and have an average particle size of about 7 µm - 15 µm. Cooling within the above-mentioned temperature range is preferably conducted such that the cooling rate becomes almost constant.

A next important process for controlling the crystal characteristics is the addition of seed crystal in (B1) in the specific temperature range. Addition of a seed crystal for crystallization of a compound is a known preferable method for precipitating crystals, as demonstrated in US4650884 with regard to citalopram hydrobromide. In this publication, a seed crystal is added when the cooling temperature is 20°C. By reproductive crystallization according to this method by the present inventors, it was revealed that suitable growth of the precipitated crystals was not achieved, with a considerable amount of fine crystals having a particle size of less than 5 µm. When fine crystals are contained in a great amount, filterability after crystallization for the preparation of a pharmaceutical bulk becomes poor, and the fine particles scatter, to which particles the workers are exposed during production of pharmaceutical preparations.

The present inventors have also found that, in (B1), cooling to the temperature range of preferably not less than 30°C and less than 48°C, more preferably not less than 40°C and less than 48°C, followed by addition of a seed crystal to allow for crystallization, results in precipitation of crystals suitably grown. Thus, the particle size of the crystals can be controlled well, and generation of fine crystals having a particle size of less than 5 µm can be further inhibited by this method.

When the temperature, at which a seed crystal is added, is less than 30°C, crystallization occurs at once and fine crystals tend to increase. When it is not less than 48°C, the seed crystal tends to dissolve easily, and the effect of the addition of the seed crystal becomes low. The amount of the seed crystal added is not subject to any particular limitation, but it is preferably 0.0001 wt% - 1 wt%, more preferably 0.001 wt% - 0.01 wt%, relative to citalopram hydrobromide used for crystallization.

In (A1) in Method 1, citalopram hydrobromide is dissolved in a specific solvent preferably by heating to 50°C - 80°C, more preferably 55°C - 75°C.

The specific solvent used in Method 1 contains at least one member selected from the group consisting of alcohol having 1 to 3 carbon atoms (e.g., methanol, ethanol, isopropyl alcohol, 1-propanol), water and acetone, in a proportion of preferably not less than 70 wt%, more preferably not less than 80 wt%. The solvent that may be contained in addition to alcohol having 1 to 3 carbon atoms, water and acetone is not subject to any particular limitation, and may be any as long as it is miscible with these solvents and is not reactive with citalopram hydrobromide. Examples thereof include toluene, xylene, tetrahydrofuran, diisopropyl ether, 1,2-dimethoxyethane, diethylene glycol dimethyl ether and the like. These may be used alone or in a combination of 2 or more solvents, wherein the boiling point thereof is preferably higher than the above-mentioned heating temperature. More preferable examples of the above-mentioned specific solvent include those containing at least one member selected from the group consisting of methanol, ethanol, isopropyl alcohol and 1-propanol in a proportion of not less than 80 wt%, particularly preferably those containing methanol and isopropyl alcohol in a proportion of not less than 90 wt%.

The amount of the above-mentioned specific solvent varies depending on the kind of each solvent used solely or in combination, the temperature at which citalopram hydrobromide is heated for dissolution and the like. However, it is generally preferably 2 L - 30 L, more preferably 4 L - 16 L, per 1 kg of citalopram hydrobromide. When a mixed solvent containing methanol and isopropyl alcohol is used, it is particularly preferable that

the amounts thereof be 2 L - 5 L of methanol and 4 L - 8L of isopropyl alcohol, both per 1 kg of citalopram hydrobromide.

According to Method 1, after the completion of step 2, the crystallization of citalopram hydrobromide is preferably completed after aging at a temperature below the final cooling temperature (preferably -10°C to 30°C, more preferably 0°C to 20°C) for preferably 30 minutes - 48 hours, more preferably 2 hours - 24 hours, to achieve an improved yield.

By the foregoing steps, the characteristics of citalopram hydrobromide crystals can be controlled. For example, to shift the particle size distribution of the crystals toward that of the crystals having a greater particle size, the cooling rate may be slowed or a seed crystal may be added in (B1). To increase the aspect ratio, the cooling rate or the stirring speed may be slowed in (B1).

Method 2

The second method for crystallizing citalopram hydrobromide according to the present invention includes the steps of:

(A2) dissolving citalopram hydrobromide in a specific solvent by heating,

(B2) crystallizing citalopram hydrobromide by cooling the obtained solution,

(C2) heating the resulting crystals to dissolve a part of the obtained crystals, and

(D2) recrystallizing citalopram hydrobromide while controlling a cooling rate

(hereinafter this method is to be referred to as Method 2). The crystals obtained in Method 2 generally contain fine crystals in a smaller proportion than in Method 1. In Method 2, the steps of (A2) to (D2) can be performed in one pot. In each step, it is preferable that the particles be stirred at 10 rpm - 250 rpm, particularly at 50 rpm - 220 rpm, so that the temperature of the system as a whole will be uniform and the crystals after stirring will not have a smaller aspect ratio.

In Method 2, the control of the cooling rate of the solution within the specific temperature range in (D2) is particularly important for the control of the crystal characteristics. That is, the present inventors have found that

the cooling rate of the solution particularly within the temperature range of from (heating temperature in (C2)) to (this heating temperature - 30°C) in (D2) in Method 2 affects the crystal particle size of the ultimate crystals. For example, the crystals obtained when the average cooling rate of the solution within this temperature range is controlled to not less than 30°C/hour and not more than 90°C/hour contain crystals having a particle size of less than 5 μm in a proportion of 35% at most, particularly not more than 25%, and have an average particle size of about 10 μm - 25 μm . The crystals obtained when it is controlled to not less than 1°C/hour and less than 30°C/hour contain crystals having a particle size of less than 5 μm in a proportion of 35% at most, particularly not more than 15%, and have an average particle size of about 15 μm - 30 μm . Cooling within the above-mentioned temperature range is preferably conducted such that the cooling rate becomes almost constant.

A next important process for controlling the crystal characteristics is the addition of a seed crystal within the specific temperature range in (B2) and dissolution by heating in (C2).

In (B2), the solution obtained in (A2) is cooled to the temperature range of preferably not less than 30°C and less than 48°C, particularly preferably not less than 40°C and less than 48°C. The present inventors have found that, in (B2), cooling to the temperature range of preferably not less than 30°C and less than 48°C, particularly preferably not less than 40°C and less than 48°C, which is followed by addition of a seed crystal to allow for crystallization results in the precipitation of crystals suitably grown. Thus, the particle size of the crystals can be controlled well, and generation of fine crystals having a particle size of less than 5 μm can be further inhibited.

When the temperature at which a seed crystal is added is less than 30°C, crystallization occurs at once and fine crystals tend to increase. When it is not less than 48°C, the seed crystal tends to dissolve easily, and the effect of the addition of the seed crystal becomes low. The amount of the seed crystal added is not subject to any particular limitation, but it is preferably 0.0001 wt% - 1 wt%, more preferably 0.001 wt% - 0.01 wt%, relative

to citalopram hydrobromide used for crystallization.

In (C2), a part of the crystals obtained in (B2) is dissolved by heating, thereby preventing complete dissolution of the obtained crystals. This has a consequence that the finally
5 obtained crystals have a greater average particle size. This is considered to be attributable to the fact that the crystals not dissolved by heating in (B2) become the seed crystal in (D2). Therefore, combined with the addition of the seed crystal in (B2), crystals having a greater average particle size as compared to
10 that obtained without such combination can be ultimately obtained. As used herein, the "part of the crystals obtained in (B2)" is not subject to any particular limitation, but it is preferably 10 wt% - 90 wt%, more preferably 20 wt% - 80 wt%, of the crystals obtained in (B2).

The temperature in (C2) is not subject to any particular
15 limitation as long as it can dissolve only a part of the crystals obtained in (B2). Note that the crystals should not be dissolved completely in (B2). It is preferable that a part of the crystals be dissolved by heating to not less than 48°C and not more than
20 60°C, particularly not less than 48°C and not more than 55°C. When this temperature is less than 48°C, the crystals cannot be dissolved to the desired degree in this step, and when it exceeds 60°C, the crystals are dissolved to the degree beyond the intended degree in this step, both causing an increase in the content of
25 the crystals having a particle size of less than 5 μ m, as compared to (C2) performed in the above-mentioned temperature range.

In Method 2, the cooling rate controlled in (B2) also affects the particle size of the final crystals, as is the case with the aforementioned control of cooling rate in (D2), only to
30 a smaller degree than in (D2). In (B2), the present inventors have found that the cooling rate within the temperature range of preferably 0°C - 80°C, particularly 20°C - 40°C, affects the crystal particle size of citalopram hydrobromide. The cooling in the above-mentioned temperature range is preferably performed
35 such that the cooling rate becomes almost the same.

In (A2) in Method 2, citalopram hydrobromide is dissolved in a specific solvent preferably by heating to 50°C - 80°C, more preferably 55°C - 75°C.

The specific solvent used in Method 2 contains at least one member selected from the group consisting of alcohol having 1 to 3 carbon atoms (e.g., methanol, ethanol, isopropyl alcohol, 1-propanol), water and acetone, preferably in a proportion of not less than 70 wt%, more preferably not less than 80 wt%. The solvent that may be contained in addition to alcohol having 1 to 3 carbon atoms, water and acetone is not subject to any particular limitation, and may be any as long as it is miscible with these solvents and is not reactive with citalopram hydrobromide.

Examples thereof include toluene, xylene, tetrahydrofuran, diisopropyl ether, 1,2-dimethoxyethane, diethylene glycol dimethyl ether and the like. These may be used alone or in a combination of 2 or more solvents, wherein the boiling point thereof is preferably higher than the above-mentioned heating temperature. More preferable examples of the above-mentioned specific solvent include those containing at least one member selected from the group consisting of methanol, ethanol, isopropyl alcohol and 1-propanol in a proportion of not less than 80 wt%. Those containing methanol and isopropyl alcohol in a proportion of not less than 90 wt% are particularly preferable.

The amount of the above-mentioned specific solvent varies depending on the kind of each solvent used solely or in combination, the temperature at which citalopram hydrobromide is heated for dissolution and the like. However, it is generally preferably 2 L - 30 L, more preferably 4 L - 16 L, per 1 kg of citalopram hydrobromide. When a mixed solvent containing methanol and isopropyl alcohol is used, it is particularly preferable that the amounts thereof be 2 L - 5 L of methanol and 4 L - 8 L of isopropyl alcohol, both per 1 kg of citalopram hydrobromide.

In Method 2, (C2) and (D2) may be sequentially repeated after (D2). The crystallization of citalopram hydrobromide is preferably completed after aging at a temperature below the final cooling temperature (preferably -10°C to 30°C, more preferably 0°C to 20°C) for preferably 30 minutes - 48 hours, more preferably 2 hours - 24 hours, to achieve an improved yield.

By the foregoing steps, the characteristics of citalopram hydrobromide crystals can be controlled. For example, to shift the particle size distribution of the crystals toward that of the

crystals having a greater particle size, the cooling rate may be slowed in (B2) and (D2), a seed crystal may be added in (B2), and the crystals may be heated for dissolution at a specific temperature in (C2). To increase the aspect ratio, the cooling rate or the stirring speed may be slowed in (B2) and (D2).

In Method 1 and Method 2, the obtained crystals are isolated and purified by conventional methods, such as washing with a suitable solvent after filtration.

The citalopram hydrobromide crystals wherein 35% at most, preferably not more than 30%, more preferably not more than 20%, thereof has a crystal particle size of less than 5 μm , and further the crystals wherein not less than 10%, more preferably not less than 20%, thereof preferably has a particle size of not less than 20 μm , can improve the filterability after crystallization for the production of pharmaceutical bulk and the like. In addition, they can improve the problems of scattering of fine particles and exposure of the workers to the fine particles during production of a pharmaceutical preparation, and the like. These crystals can be obtained by each method as follows.

In Method 1, the average cooling rate of the solution within the temperature range of from 20°C to 40°C in (B1) is controlled to not less than 0.5°C/hour and not more than 60°C/hour, preferably by adding a seed crystal at a temperature range of not less than 30°C and less than 48°C in (B1).

In Method 2, the cooling rate of the solution in the temperature range of from ((heating temperature in (C2)) to (this heating temperature - 30°C) in (D2) is controlled to not less than 1°C/hour and not more than 90°C/hour, or preferably a seed crystal is added at a temperature range of not less than 30°C and less than 48°C in (B2), or a part of the obtained crystals in (B2) is heated for dissolution in (C2).

The citalopram hydrobromide crystals having an average aspect ratio of generally not less than 2.0, preferably not less than 3, and more preferably not less than 4.5, and generally not more than 9.0, preferably not more than 7, and more preferably not more than 6.0, can improve the problems such as poor filtering performance after crystallization for the preparation of a pharmaceutical bulk and poor fluidity when crystals are being

taken out, and can be advantageously used for the production of citalopram hydrobromide and the production of pharmaceutical products. From this viewpoint, the crystals having an average aspect ratio of not less than 4.5 and not more than 6.0 are particularly preferable. These crystals can be obtained by each method as follows.

In Method 1, the average cooling rate of the solution in the temperature range of 20°C - 40°C in (B1) is controlled to not less than 0.5°C/hour and not more than 60°C/hour, and preferably the stirring speed is set for 10 rpm - 250 rpm in (B1). In (B1), the speed at the tip of an agitation part is generally set to 0.1 m/s - 2.5 m/s, preferably 0.5 m/s - 2.0 m/s.

In Method 2, the cooling rate of the solution within the temperature range of from (heating temperature in (C2)) to (this heating temperature - 30°C) is controlled to not less than 1°C/hour and not more than 90°C/hour in (D2), and preferably the stirring speed is set for 10 rpm - 250 rpm in (B2) and (D2). In (B2) and (D2), the speed at the tip of an agitation part is generally set to 0.1 m/s - 2.5 m/s, preferably 0.5 m/s - 2.0 m/s.

By setting the speed at the tip of an agitation part to generally from not less than 2.5 m/s to less than 5.0 m/s, preferably from not less than 3.0 m/s to not more than 4.0 m/s, in (B1) in Method 1 and (B2) and (D2) in Method 2, crystals having an average aspect ratio of not less than 2.5 and less than 4.5 can be obtained. The crystals having an average aspect ratio of not less than 2.5 and less than 4.5 are preferable because they can be used as they are as a pharmaceutical preparation.

The citalopram hydrobromide to be used may be any that can be obtained by any method. It is preferably purified by active charcoal and the like before being subjected to the crystallization according to the present invention. The citalopram hydrobromide can be obtained, for example, according to the disclosure in, for example, US4650884.

The present invention is explained in more detail in the following by referring to Examples that do not limit the present invention in any way.

The powder X-ray crystal diffraction and average particle

size were measured in the following Reference Example, Examples and Comparative Example, according to the following methods.

Powder X-ray crystal diffraction: Rigaku Corporation

MiniFlex

Irradiated X-ray CuK α

Average particle size: SHIMADZU CORPORATION SALD1100

(medium : diisopropyl ether)

Reference Example 1

Citalopram base (167 g) was dissolved in acetone (837 ml).

Thereeto was blown in a hydrobromic acid gas at 15°C - 40°C to make the pH of the solution 3. When about 1/2 of the hydrobromic acid gas was blown in, crystals began to precipitate. The reaction mixture was stirred at 25°C - 35°C for 1 hour, and at 0°C - 5°C for 1 hour. The resulting crystals were filtered, washed with cold acetone (167 ml) and dried in vacuo at 50°C - 70°C (5 Torr - 30 Torr) for 3 hours to give citalopram hydrobromide crystals (162 g). The crystals had an HPLC purity of 97.3% (254 nm).

melting point 180.3°C

¹H-NMR (DMSO-d₆, 400 MHz): shown in Figure 1.

Powder X-ray crystal diffraction pattern: shown in Figure 2.

Example 1

Citalopram hydrobromide (5.0 g) was synthesized according to Reference Example 1 and dissolved in methanol (15 ml) at 60°C. Thereeto was added active charcoal (0.5 g) and the mixture was stirred for 15 minutes. The active charcoal was filtered off and isopropyl alcohol (25 ml) was added. The mixture was heated to 58°C to dissolve the crystals. When the mixture was cooled to 43°C while stirring at 200 rpm (propeller diameter: 0.05 m, speed at the tip of propeller: $0.05 \times (200/6) \times 3.14 = \text{ca. } 0.5 \text{ m/s}$), the seed crystal (about 0.1 mg) of citalopram hydrobromide was added. The mixture was cooled to 40°C over 15 minutes and then cooled from 40°C to 20°C over 5 hours while stirring at 200 rpm to make the cooling rate almost the same (average cooling rate: 4°C/hour). The cooled suspension was stirred at 20°C for 18 hours (retention time at 20°C) at 200 rpm and the resulting crystals were filtrated. The crystals were dried at 50°C (0.67 kPa - 1.33 kPa) for 3 hours in vacuo while stirring them at 30 rpm, and stood for drying at 70°C (0.4 kPa - 0.8 kPa) for 15 hours.

The obtained citalopram hydrobromide crystals weighed 4.2 g, and the crystals had a melting point of 184.4°C and an HPLC purity of 99.9% (254 nm).

powder X-ray crystal diffraction pattern: shown in Figure 3
crystal particle size distribution

less than 5 μm 17.1%,
not less than 5 μm and less than 10 μm 22.2%,
not less than 10 μm and less than 20 μm 29.1%,
not less than 20 μm and less than 40 μm 27.0%,
not less than 40 μm 4.6% (shown in Figure 6)

average particle size 12.9 μm

micrograph (400 magnifications): shown in Figure 13

average aspect ratio: 4.8

Examples 2 - 3

The procedure followed Example 1 except that the cooling time of from 40°C to 20°C and the retention time at 20°C were changed to those shown in Table 1.

Table 1

Ex.	Cooling time of from 40°C to 20°C (cooling rate in parenthesis)	Retention time at 20°C	Crystal particle size distribution (D)	Average particle size (μm)
2	30 min (40°C/hour)	19 hr 30 min	D<5 μm 31.4% 5 μm \leq D<10 μm 30.3% 10 μm \leq D<20 μm 25.7% 20 μm \leq D<40 μm 12.2% D \geq 40 μm 0.3% (shown in Figure 7)	7.8
3	1 hr (20°C/hour)	19 hr	D<5 μm 19.6% 5 μm \leq D<10 μm 26.0% 10 μm \leq D<20 μm 31.0% 20 μm \leq D<40 μm 21.6% D \geq 40 μm 1.6% (shown in Figure 8)	11.0

Example 4

Citalopram hydrobromide (5.0 g) was synthesized according to Reference Example 1 and dissolved in methanol (15 ml) at 50°C. Thereto was added active charcoal (0.5 g) and the mixture was stirred for 15 minutes. The active charcoal was filtered off and isopropyl alcohol (25 ml) was added. The mixture was heated to 58°C to dissolve the crystals. When the mixture was cooled to 43°C while stirring at 200 rpm (propeller diameter: 0.05 m), the seed crystal (0.1 mg) of citalopram hydrobromide was added. The mixture was cooled to 41°C over 15 minutes and heated to 50°C over 5 minutes, and then stirred at the same temperature for 80 minutes to allow dissolution of a part of the crystals (about half). Then, the mixture was cooled from 50°C to 20°C over 2.5 hours to make the cooling rate almost the same (average cooling rate: 12°C/hour). The cooled suspension was stirred at 20°C for 17.5 hours at 200 rpm and the precipitated crystals were filtrated. The crystals were dried at 50°C (0.67 kPa - 1.33 kPa) for 3 hours in vacuo with stirring at 30 rpm, and stood for drying at 70°C (0.4 kPa - 0.8 kPa) for 15 hours.

The obtained citalopram hydrobromide crystals weighed 4.1 g and the crystals had an HPLC purity of 99.9% (254 nm) and a melting point of 184.7°C.

powder X-ray crystal diffraction pattern: shown in Figure 4.
crystal particle size distribution

less than 5 μm 9.6%,
not less than 5 μm and less than 10 μm 12.3%,
not less than 10 μm and less than 20 μm 20.9%,
not less than 20 μm and less than 40 μm 31.7%,
not less than 40 μm 25.5% (shown in Figure 9)

average particle size 23.8 μm
micrograph (400 magnifications): shown in Figure 14
average aspect ratio: 5.4

Example 5

The procedure followed Example 4 except that the cooling time of from 50°C to 20°C and the retention time at 20°C were changed to those shown in Table 2.

Table 2

Ex.	Cooling time of from 50°C to 20°C (cooling rate in parenthesis)	Retention time at 20°C	Crystal particle size distribution (D)	Average particle size (μm)
5	30 min (60°C/hr)	19 hr 30 min	D<5 μm 15.6% 5 μm \leq D<10 μm 19.3% 10 μm \leq D<20 μm 26.9% 20 μm \leq D<40 μm 31.0% D \geq 40 μm 7.2% (shown in Figure 10)	15.1

Example 6

5 Citalopram hydrobromide (18.2 kg) was synthesized according to Reference Example 1 and dissolved in methanol (37.4 kg) at 45°C - 50°C. Thereto was added active charcoal (1.8 kg) and the mixture was stirred for 55 minutes. The active charcoal was filtered off and washed with warm methanol (7.2 kg). The washings and the filtrate were combined and heated to about 60°C. Isopropyl alcohol (71.9 kg) heated to 40°C - 50°C was added, and upon confirmation of dissolution at 62°C, the mixture was cooled with stirring at 169 rpm (diameter of impeller of agitator (Pfaudler type): 0.45 m, speed at the tip of impeller of agitator: 10 ca. 4.0 m/s). When the mixture was cooled to 44°C, the seed crystal (about 400 mg) of citalopram hydrobromide was added. Stirring at 40°C - 43°C for 30 minutes enabled confirmation of the precipitation of the crystals at 41.5°C.

The mixture was cooled from 40°C to 20°C over 4 hours (average cooling rate: 5°C/hour), and further cooled to 10°C over 20 3 hours. The suspension was stirred at 5°C - 10°C at 80 rpm for 13 hours and the crystals were filtrated. A mixture of isopropyl alcohol (9.6 kg) and methanol (4.8 kg) was cooled to 0°C - 10°C. The crystals were washed with this mixture. The wet crystals 25 (19.8 kg) were dried under reduced pressure (10 kPa - 1.33 kPa), at 30°C - 50°C for 10 hours, at 45°C - 55°C for 5 hours, and at 70°C - 75°C for 8 hours, with stirring at 30 rpm.

The obtained citalopram hydrobromide crystals weighed 15.0 kg (recrystallization yield : 82%) and the crystals had an HPLC 30 purity of 99.75% (220 nm) and a melting point of 185.0°C.

crystal particle size distribution

less than 5 μm 12.6%,

not less than 5 μm and less than 10 μm 20.2%,

not less than 10 μm and less than 20 μm 35.5%,

not less than 20 μm and less than 40 μm 29.3%,

not less than 40 μm 2.4% (shown in Figure 11)

average particle size: 14.6 μm

micrograph (400 magnifications): shown in Figure 15.

average aspect ratio: 3.0

bulk density: 0.2 g/ml

tap density 0.42 g/ml

Comparative Example 1

Citalopram hydrobromide (5.0 g) was synthesized according to Reference Example 1 and dissolved in methanol (15 ml) at 50°C. Thereto was added active charcoal (0.5 g) and the mixture was stirred for 15 minutes. The active charcoal was filtered off and isopropyl alcohol (25 ml) was added. The mixture was heated to 58°C to dissolve the crystals. The mixture was cooled to 40°C over 5 min and then cooled from 40°C to 20°C over 5 min with stirring at 200 rpm. When the mixture was cooled to 20°C, the seed crystal (0.1 mg) was added. The mixture was stirred at the same temperature for 20 hours and the crystals were filtrated. The crystals were dried at 50°C (0.67 kPa - 1.33 kPa) for 3 hours in vacuo with stirring at 30 rpm, and stood for drying at 70°C (0.4 kPa - 0.8 kPa) for 15 hours.

The obtained citalopram hydrobromide crystals weighed 4.3 g and the crystals had an HPLC purity of 99.9% (254 nm) and a melting point of 184.2°C.

powder X-ray crystal diffraction pattern: shown in Figure 5
crystal particle size distribution

less than 5 μm 41.9%,

not less than 5 μm and less than 10 μm 30.2%,

not less than 10 μm and less than 20 μm 23.9%,

not less than 20 μm and less than 40 μm 4.0%,

not less than 40 μm 0% (shown in Figure 12)

average particle size: 6.3 μm

micrograph (400 magnifications): shown in Figure 16

According to the method of the present invention, the crystal characteristics of citalopram hydrobromide, such as particle size, particle size distribution, aspect ratio and the like, can be controlled easily and industrially. In addition,
5 citalopram hydrobromide crystals having crystal characteristics useful as a pharmaceutical bulk can be provided.

This application is based on application No. 2000-133995 filed in Japan, the contents of which are incorporated hereinto by
10 reference.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
222